SEVERE ALLERGY CARE PLAN AND MEDICATION AUTHORIZATION FORM

	First	***	Birthdate:
Last	First	M.I.	
chool Name:	School Year:	Grade:	
tep 1: Identification of all	ergen- This section to be com	pleted by Medical Prov	ider.
	Previous:		
levere	Symptoms		
llergy to:	(if known):		
tep 2: Treatment Protoco	I- This section to be completed	l by Medical Provider o	nly.
Severe Symptoms If any of the following severe symptoms are noted LUNG: Short of breath, wheeze, repetitive cough HEART: Pale, blue, faint, weak pulse, dizzy THROAT: Tight, hoarse, trouble breathing/swallowing MOUTH: Significant swelling of the tongue and/or lips.		1. INJECT EPINEPHRINE IMMEDIATELY	
		2. CALL 911	
		3. GIVE ADDITIONAL MEDICATIONS (IF ORDERED BY PHYSICIAN)	
SKIN: Many hives over b	ody, widespread redness		and raise legs. If breathing is
GUT: Repetitive vomiting	g or severe diarrhea	difficult or they are	vomiting, let them sit up or lie o
OTHER: Feeling something bad is about to happen, Confusion, anxiety		their side. For insect stings/bites only: remove stinger if present.	
Mild	Symptoms	1 GIVE medical	tion indicated. See below.
If the following symptoms are	e noted, give medication		
indicated. (Orders below)		2. Stay with student	t, alert emergency contacts.
THROAT: hoarse, persistent or	ough	3. Watch student cl	osely for changes. If symptoms
MOUTH: Itchy or tingling mou		worsen, or severe s	ymptoms appear, GIVE
SKIN: A few hives/rash, mild it	tch	EPINEPHRINE and	refer to treatment protocol
GUT: Mild nausea/discomfort	☐ Epi-pen ☐ Antihistamine	above for severe s	umntome
	C chi-heii C vanishinini		
OTHER:		4. If an epi-pen is a	administered, call 911.
If checked, give epinephrine If checked, give epinephrine	☐ Epi-pen ☐ Antihistamine immediately for ANY symptoms if the simmediately if the student was definite tions-This section to be complete.	4. If an epi-pen is a student was likely exposed to by exposed to the allergen/sti- eted Medical Provider of	the allergen/sting. ng, even if no symptoms are no
If checked, give epinephrine If checked, give epinephrine	☐ Epi-pen ☐ Antihistamine immediately for ANY symptoms if the s immediately if the student was definite	4. If an epi-pen is a student was likely exposed to by exposed to the allergen/sti	the allergen/sting. ng, even if no symptoms are no only. Bronchodilator
If checked, give epinephrine If checked, give epinephrine Step 3: Authorized Medica Name of Medication	☐ Epi-pen ☐ Antihistamine immediately for ANY symptoms if the simmediately if the student was definite tions-This section to be compl Epinephrine Auto injector	4. If an epi-pen is a student was likely exposed to by exposed to the allergen/sti eted Medical Provider of Antihistamine	the allergen/sting. ng, even if no symptoms are no
If checked, give epinephrine If checked, give epinephrine Itep 3: Authorized Medica Name of Medication	☐ Epi-pen ☐ Antihistamine immediately for ANY symptoms if the simmediately if the student was definite tions-This section to be compl Epinephrine Auto injector	4. If an epi-pen is a student was likely exposed to by exposed to the allergen/sti eted Medical Provider of Antihistamine	the allergen/sting. ng, even if no symptoms are no only. Bronchodilator
If checked, give epinephrine If checked, give epinephrine Itep 3: Authorized Medica Name of Medication Purpose	☐ Epi-pen ☐ Antihistamine immediately for ANY symptoms if the simmediately if the student was definite tions-This section to be compl Epinephrine Auto injector	4. If an epi-pen is a student was likely exposed to by exposed to the allergen/sti eted Medical Provider of Antihistamine	the allergen/sting. ng, even if no symptoms are no only. Bronchodilator
If checked, give epinephrine If checked, give epinephrine Step 3: Authorized Medica Name of Medication Purpose Strength	□ Epi-pen □ Antihistamine immediately for ANY symptoms if the simmediately if the student was definite tions-This section to be compl Epinephrine Auto injector 1.	4. If an epi-pen is a student was likely exposed to by exposed to the allergen/sti eted Medical Provider of Antihistamine	the allergen/sting. ng, even if no symptoms are no only. Bronchodilator
If checked, give epinephrine If checked, give epinephrine Itep 3: Authorized Medical Name of Medication Purpose Strength Medication Form	□ Epi-pen □ Antihistamine immediately for ANY symptoms if the s immediately if the student was definite tions- This section to be compl Epinephrine Auto injector 1. □ 0.3mg □ 0.15mg Auto-injector Injected intramuscularly into	4. If an epi-pen is a student was likely exposed to by exposed to the allergen/sti eted Medical Provider of Antihistamine	the allergen/sting. ng, even if no symptoms are no only. Bronchodilator
If checked, give epinephrine If checked, give epinephrine Itep 3: Authorized Medical Name of Medication Purpose Strength Medication Form Route of Admin	□ Epi-pen □ Antihistamine immediately for ANY symptoms if the s immediately if the student was definite tions- This section to be compl Epinephrine Auto injector 1. □ 0.3mg □ 0.15mg Auto-injector	4. If an epi-pen is a student was likely exposed to by exposed to the allergen/sti eted Medical Provider of Antihistamine	the allergen/sting. ng, even if no symptoms are no only. Bronchodilator
If checked, give epinephrine If checked, give epinephrine Itep 3: Authorized Medical Name of Medication Purpose Strength Medication Form Route of Admin Or frequency if PRN	□ Epi-pen □ Antihistamine immediately for ANY symptoms if the s immediately if the student was definite tions- This section to be compl Epinephrine Auto injector 1. □ 0.3mg □ 0.15mg Auto-injector Injected intramuscularly into	4. If an epi-pen is a student was likely exposed to by exposed to the allergen/sti eted Medical Provider of Antihistamine	the allergen/sting. ng, even if no symptoms are no only. Bronchodilator
If checked, give epinephrine If checked, give epinephrine Itep 3: Authorized Medica Name of Medication Purpose Strength Medication Form Route of Admin Or frequency if PRN Precautions, instructions,	□ Epi-pen □ Antihistamine immediately for ANY symptoms if the s immediately if the student was definite tions- This section to be compl Epinephrine Auto injector 1. □ 0.3mg □ 0.15mg Auto-injector Injected intramuscularly into	4. If an epi-pen is a student was likely exposed to by exposed to the allergen/sti eted Medical Provider of Antihistamine	the allergen/sting. ng, even if no symptoms are no only. Bronchodilator
If checked, give epinephrine If checked, give epinephrine If checked, give epinephrine Itep 3: Authorized Medica Name of Medication Purpose Strength Medication Form Route of Admin Or frequency if PRN Precautions, instructions, Adverse effects or comments	□ Epi-pen □ Antihistamine immediately for ANY symptoms if the s immediately if the student was definite tions- This section to be compl Epinephrine Auto injector 1. □ 0.3mg □ 0.15mg Auto-injector Injected intramuscularly into	4. If an epi-pen is a student was likely exposed to by exposed to the allergen/sti eted Medical Provider of Antihistamine	the allergen/sting. ng, even if no symptoms are no only. Bronchodilator
If checked, give epinephrine If checked, give epinephrine If checked, give epinephrine Itep 3: Authorized Medical Name of Medication Purpose Strength Medication Form Route of Admin Or frequency if PRN Precautions, instructions, Adverse effects or comments Can the student carry and self-	□ Epi-pen □ Antihistamine immediately for ANY symptoms if the simmediately if the student was definite tions- This section to be compl Epinephrine Auto injector 1. □ 0.3mg □ 0.15mg Auto-injector Injected intramuscularly into lateral thigh	4. If an epi-pen is a student was likely exposed to the allergen/still eted Medical Provider (Antihistamine 2.	the allergen/sting. ng, even if no symptoms are no only. Bronchodilator 3.
If checked, give epinephrine If checked, give	□ Epi-pen □ Antihistamine immediately for ANY symptoms if the s immediately if the student was definite tions- This section to be compl Epinephrine Auto injector 1. □ 0.3mg □ 0.15mg Auto-injector Injected intramuscularly into lateral thigh	4. If an epi-pen is a student was likely exposed to by exposed to the allergen/stilleted Medical Provider (Antihistamine 2.	the allergen/sting. ng. even if no symptoms are no only. Bronchodilator 3.
If checked, give epinephrine If checked, give	□ Epi-pen □ Antihistamine immediately for ANY symptoms if the simmediately if the student was definite tions- This section to be compl Epinephrine Auto injector 1. □ 0.3mg □ 0.15mg Auto-injector Injected intramuscularly into lateral thigh □ Yes □ No The As the Medical Provider of the above ations be available for administration du	4. If an epi-pen is a student was likely exposed to be allergen/stileted Medical Provider of Antihistamine 2.	the allergen/sting. ng. even if no symptoms are no only. Bronchodilator 3.
If checked, give epinephrine If checked, give	□ Epi-pen □ Antihistamine immediately for ANY symptoms if the simmediately if the student was definite tions- This section to be compl Epinephrine Auto injector 1. □ 0.3mg □ 0.15mg Auto-injector Injected intramuscularly into lateral thigh □ Yes □ No The As the Medical Provider of the above ations be available for administration du	4. If an epi-pen is a student was likely exposed to be allergen/stileted Medical Provider of Antihistamine 2.	the allergen/sting. ng. even if no symptoms are no only. Bronchodilator 3.

Step 4: Parent/Guardian to complete Cell phone: Relationship: Other phone: Emergency Contacts: Office phone number: Child's Physician Name: If a medication must be taken during the school day or during a school sponsored overnight trip, it is necessary, in accordance with California Education Code Section 49423, to have a written statement on file. The statement must be signed by the parent/guardian and the physician indicating a desire that designated school personnel assist the student with medication administration. The authorization must be made annually and/or whenever a change occurs. Education Code requires that ALL medications, prescription and over-the-counter must have a completed statement from BOTH the physician AND parent/guardian BEFORE they can be administered. Medication must be provided in the original container labeled with student's name, medication name, dose/strength and specific administration directions. Parent/Guardian Authorization: As the parent/guardian of the above named child, I request that designated school personnel assist in the administration of medication prescribed by the Medical Provider. I give consent for the Medical Provider and designated school personnel to communicate directly, regarding the administration of the medication. I understand it is my responsibility to bring all medication safely to the school and I agree to refill or replace medication as necessary. I understand that the medication will be stored in a locked area unless the Medical Provider indicates that my child is capable of carrying and self-administering it. I hereby release the school district and all school personnel from civil liability if my child suffers an adverse reaction as the result of self-administering prescription auto-injectable epinephrine or prescription inhaled asthma medication. Name of Parent/Guardian: Signature of Parent/Guardian: Date: Student Statement (required if authorized to self-carry medication) I understand that I am allowed to carry and self-administer ONLY the medications listed above. I agree to use the medication as instructed by my physician and not to share with other people. I understand that if I share the medication with others, I will be held accountable for my actions and that I will face disciplinary action. Signature of Student: _ Date: How to administer Epi-Pen autoinjector Swing and firmly push Remove the orange tip against outer thigh so it blue safety "clicks" release by pulling straight up without AND HOLD on thigh bending or approx. 10 seconds twisting it to deliver drug